

Comment on the possible electron neutrino excess in the Super-Kamiokande atmospheric neutrino experiment

C. W. Kim

*Department of Physics and Astronomy, The Johns Hopkins University, Baltimore, MD21218,
USA*

*Korea Institute for Advanced Study, 207-43 Cheongryangri-dong, Dongdaemun-gu, Seoul 130-012,
Korea*

U. W. Lee

Department of Physics, Mokpo National University, Muan-gun, Chonnam 534-729, Korea

Email address: cwkim@kias.kaist.ac.kr, leeuw@chungkye.mokpo.ac.kr

Abstract

We investigate the implications of a possible excess of the electron neutrino events in the Super-Kamiokande atmospheric neutrino experiment. The excess, if real, leads to constraints on the $\nu_\mu \leftrightarrow \nu_e$, $\nu_\mu \leftrightarrow \nu_\tau$ and $\nu_e \leftrightarrow \nu_\tau$ oscillation parameters. Specifically, the excess implies that $\nu_e - \nu_\mu$ and $\nu_\mu - \nu_\tau$ mixings are large whereas $\nu_e - \nu_\tau$ mixing is small. We show that the electron neutrino excess favors the large mixing angle MSW solution of the solar neutrino problem.

PACS number(s): 14.60Lm, 14.60.Pq, 25.65

Recently, the Super-Kamiokande Collaboration has reported the evidence for the neutrino oscillation from the atmospheric neutrino experiment [1]. It has been remarkably well demonstrated that the data are consistent with the $\nu_\mu \leftrightarrow \nu_\tau$ oscillations with $\sin^2 2\theta > 0.82$ and $5 \times 10^{-4} < \Delta m^2 < 6 \times 10^{-3} \text{eV}^2$ at 90 % confidence level. However, the data appear to imply that there exists an excess of the number of detected electrons compared to the calculated number by Monte Carlo method [2–4] for the up-going low energy neutrinos. The Fig.1 in Ref. [1] is the plot of the up-down asymmetry for the e-like and μ -like events. One can easily see the electron event excess compared to the Monte Carlo calculation for the low energy electrons. The Fig.3 in Ref. [1] is the plot of the zenith angle distribution of the e-like and μ -like events for the sub-GeV and multi-GeV data sets. For the e-like events there are several data points which do not agree with the Monte Carlo calculation. In particular, a data point for the e-like events for $p < 0.4 \text{ GeV}$ with $-1.0 \leq \cos \theta \leq -0.6$ is more than several σ 's above the expected number. The Fig.4 in Ref. [1] provides another hint of the electron excess. Even though the data are well fitted with the dashed line (*i. e.* consistent with $\nu_\mu \leftrightarrow \nu_\tau$ and no ν_e oscillations), the largest L/E data of the e-like events has the largest deviation from the dashed line. Therefore, the three figures in Ref. [1] consistently suggest the existence of the electron neutrino excess for the up-going low energy neutrinos. Although there exist large uncertainties in the number of detected neutrinos calculated by Monte Carlo method [2–4], it is worthwhile to investigate possible implications of this electron neutrino excess. From the study of the excess of electrons in the framework of three-generation neutrino oscillations, we have obtained the constraints(or information) on the neutrino oscillation parameters.

For the sake of simplicity, we assume the CP conservation in the lepton sector and the relation

$$P(\nu_\alpha \rightarrow \nu_\beta) = P(\nu_\beta \rightarrow \nu_\alpha) = P_{\alpha\beta} , \quad (1)$$

where α, β represent e, μ and τ . Furthermore, we use the notation N_α^{MC} for the expected number of ν_α induced events calculated by the Monte Carlo method for the Super-

Kamiokande experiment and the N_α^{DATA} for the observed number of the ν_α induced events in the Super-Kamiokande experiment as published in Ref.[1], respectively.

With these conventions, the numbers of the detected e-like and μ -like events are given by

$$N_e^{DATA} = N_e^{MC} + [N_\mu^{MC} - N_e^{MC}] P_{e\mu} - N_e^{MC} P_{e\tau} , \quad (2)$$

$$N_\mu^{DATA} = N_\mu^{MC} - [N_\mu^{MC} - N_e^{MC}] P_{e\mu} - N_\mu^{MC} P_{\mu\tau} . \quad (3)$$

$P_{\alpha\beta}$ is the $\nu_\alpha \leftrightarrow \nu_\beta$ transition probability and $P_{\alpha\beta} \propto \sin^2 2\theta_{\alpha\beta}$ where $\theta_{\alpha\beta}$ represents an “effective” $\nu_\alpha - \nu_\beta$ mixing angle. (We caution the reader that this angle is not one of the CKM mixing angles.) As pointed out already, the Super-Kamiokande data show the existence of an excess of the up-going low energy electrons. Therefore, from Eq.(2) we have the following inequality

$$[N_\mu^{MC} - N_e^{MC}] P_{e\mu} - N_e^{MC} P_{e\tau} > 0. \quad (4)$$

If the excess of the e-like events is real, we obtain a very strong constraint on the neutrino oscillation probabilities $P_{e\mu}$ and $P_{e\tau}$ in the parameter regions where the electron neutrino excess has been detected. Equation (4) can be rewritten as

$$P_{e\mu} > \frac{1}{N_\mu^{MC}/N_e^{MC} - 1} P_{e\tau} . \quad (5)$$

This inequality is a function of N_μ^{MC}/N_e^{MC} . Although N_μ^{MC} and N_e^{MC} themselves have large uncertainties, their ratio is known to the accuracy of about 5 % even among several different flux estimates. (The uncertainty in each calculation is much smaller than 5 %.) Therefore, Eq.(5) is insensitive to theoretical errors in each neutrino flux. Let us examine the sub-GeV atmospheric neutrino data used in the recent super-Kamiokande atmospheric neutrino experiment. From Fig.1 of Ref. [1] we obtain the expected numbers of the electrons and muons as

$$\frac{N_\mu^{MC}}{N_e^{MC}} = 1.224 \pm 0.007, \quad (6)$$

for the events in the bin with $-1 < \cos \theta < -0.6$ and $p < 0.4$ GeV. Therefore, the following analysis is based on these events, i.e. All the limits on and values of the transition probabilities are valid for those neutrinos although the results on the mixing angles are of course valid in general. We emphasize here that in obtaining Eq.(6), N_μ^{MC} and N_e^{MC} must be shifted simultaneously up or down within the error bars rather than taking, *e.g.*, an upper value of N_μ^{MC} and a lower value of N_e^{MC} in order to obtain an upper bound on the ratio N_μ^{MC}/N_e^{MC} and so on. (Also, it is to be noted that the number in Eq.(6) is based on the calculation in Ref. [2] and can be slightly different for different calculations within errors of about 5 %.) Hence, with Eq.(5), we find

$$P_{e\mu} > 4.35 P_{e\tau} \quad (7)$$

for the up-going low energy neutrinos. This equation suggests that since $P_{e\mu} \leq 1$

$$P_{e\tau} < 0.23 \quad (8)$$

in the parameter regions where the electron neutrino excess was detected in the Super-Kamiokande experiment.

Now, let us calculate some bounds on $P_{e\mu}$ for the up-going low energy neutrinos. Since $P_{e\tau} \geq 0$, we have, from Eq.(2), the relation

$$P_{e\mu} \geq \frac{N_e^{DATA}/N_e^{MC} - 1}{N_\mu^{MC}/N_e^{MC} - 1} . \quad (9)$$

As mentioned already, although the ratio (N_μ^{MC}/N_e^{MC}) is known to a fairly good accuracy, the ratio (N_e^{DATA}/N_e^{MC}) has the uncertainties inherent in N_e^{MC} . If we simply use the ratio of the two central values of N_e^{DATA} and N_e^{MC} , which is 1.45, we obtain from Eq.(9), $P_{e\mu} > 2$, which is unphysical. Even after taking all the uncertainties as quoted in Fig.(3) of Ref. [1], we still obtain $P_{e\mu} > 1$, suggesting either too high N_e^{DATA} or too low N_e^{MC} (or both). This problem persists even if a somewhat higher value of N_e^{MC} as given in Ref. [3] is used. Therefore, as was done in Ref. [1] to find the best fit, we raise simultaneously N_e^{MC} and N_μ^{MC} by 15.8%. Then taking all the uncertainties into account, we find

$$P_{e\mu} \geq 0.51 . \quad (10)$$

Defining “effective” mixing angle $\theta_{e\mu}$ and “effective” $\Delta m_{e\mu}^2$ in the form of two generation neutrinos as

$$P_{e\mu} = \sin^2 2\theta_{e\mu} \sin^2 \left(1.27 \frac{\Delta m_{e\mu}^2}{\text{eV}^2} \frac{L/\text{km}}{E/\text{GeV}} \right) , \quad (11)$$

we obtain from Eq.(10)

$$\sin^2 2\theta_{e\mu} \geq 0.51 . \quad (12)$$

One can also consider an extreme rescaling by 20 % as was done in Ref. [7]. In this case we find

$$\sin^2 2\theta_{e\mu} \geq 0.33. \quad (13)$$

Thus, our result that $\sin^2 2\theta_{e\mu}$ is “significantly” larger than $10^{-3} \sim 10^{-2}$ (which is the value of the small mixing angle MSW solution) is rather robust. We, therefore, conclude that the large mixing angle MSW solution of the solar neutrino problem is favored over the small mixing angle solution.

It is interesting to note that the electron excess is prominent only for the up-going low energy ν_e -induced events. This means that these neutrinos with $p \simeq E < 0.4$ GeV have the longest distance to travel through the earth. They correspond to a data point at slightly above $(L/\text{km})/(E/\text{GeV}) = 10^4$ in Fig.4 in Ref. [1], which is indeed elevated. This implies that for other electron neutrinos, the values of $\Delta m_{e\mu}^2 L/E$ are expected to be less than $\pi/2$ (or $\sin^2 \left(1.27 \frac{\Delta m_{e\mu}^2}{\text{eV}^2} \frac{L/\text{km}}{E/\text{GeV}} \right) \ll 1$) so that $\nu_e \leftrightarrow \nu_\mu$ oscillations would not take place. For this situation to be realized, $1.27 \Delta m_{e\mu}^2 L/E$ must become roughly $\pi/2$ only when $L \simeq 2R_E$ and $E < 0.4$ GeV (we take 0.3 GeV for events with $E < 0.4$ GeV). We then find

$$\Delta m_{e\mu}^2 \simeq 10^{-5} \text{ eV}^2 , \quad (14)$$

which is indeed consistent with the value of $\Delta m_{e\mu}^2$ obtained as the MSW solution of the solar neutrino problem [5,6], further rendering a support to our conclusion. Furthermore,

the oscillation parameters are consistent with the constraint obtained from the CHOOZ experiment [8] because the “effective” $\Delta m_{e\mu}^2$ is well below the forbidden region of the CHOOZ experiment.

Next we consider $P_{\mu\tau}$. From unitarity and Eq.(10), we find

$$P_{\mu\tau} \leq 0.49 . \quad (15)$$

This does not necessarily mean that $\sin^2 2\theta_{\mu\tau} \leq 0.49$, which disagrees with $\sin^2 2\theta_{\mu\tau} \simeq 1$ implied by the two generation fit in Ref. [1]. If we adapt $\Delta m_{\mu\tau}^2 \simeq 10^{-3}\text{eV}^2$, which is the result given in Ref. [1], $\Delta m_{\mu\tau}^2 L/E$ becomes much larger than the oscillation length for $L \simeq 2R_E$ and $E < 0.4$ GeV. Thus we expect the oscillation to be in the rapid oscillation stage with the average $\left\langle \sin^2 \left(1.27 \frac{\Delta m_{\mu\tau}^2}{\text{eV}^2} \frac{L/\text{km}}{E/\text{GeV}} \right) \right\rangle = \frac{1}{2}$, so that Eq.(15) gives

$$\sin^2 2\theta_{\mu\tau} \leq 0.98 , \quad (16)$$

which is not inconsistent with the result, $\sin^2 2\theta_{\mu\tau} \simeq 1$, in Ref. [1]

In conclusion, we need the combined analysis of $\nu_e \leftrightarrow \nu_\mu$, $\nu_e \leftrightarrow \nu_\tau$ and $\nu_\mu \leftrightarrow \nu_\tau$ oscillations to explain the atmospheric neutrino data [9]. The small $\Delta m_{e\mu}^2$ of order 10^{-5}eV^2 with a large $\nu_e - \nu_\mu$ mixing angle provides an explanation for the excess of the up-going low energy e-like events. Therefore, *the large mixing angle MSW solution of the solar neutrino problem is favored over the small mixing angle solution (which is statistically favored) if the excess of the electron neutrino in the Super-Kamiokande atmospheric neutrino experiment turns out to be real.* It remains to be seen with great interest whether the electron excess persists or not in the future experiment. A large $\nu_\mu - \nu_\tau$ mixing angle with $\Delta m_{\mu\tau}^2 \simeq 10^{-3}\text{eV}^2$ is also necessary to explain the depletion of the μ -like events. The nonzero value of $P_{e\tau}$ gives the depletion of the e-like events. Since we do not see any depletion of the e-like events, we can conclude that $P_{e\tau} \simeq 0$, consistent with Eq.(8). Our findings are in agreement with the conclusion favoring the large mixing angle MSW solution of the solar problem in Ref. [10] and the last two papers in Ref[9]. A detail numerical analysis with proper statistical considerations of our study will be presented elsewhere.

ACKNOWLEDGMENTS

U. W. Lee gratefully thanks KIAS for the hospitality during his visiting and this work is supported in part by Basic Science Research Institute of Mokpo National University. The authors would like thank Carlo Giunti for very helpful comments and suggestions.

REFERENCES

- [1] Super-Kamiokande Collaboration, Y. Fukuda, *et al.*, Phys. Rev. Lett. 81, 1562(1998).
- [2] M. Honda, *et. al.*, Phys. Lett. B248, 193(1990); M. Honda, *et. al.*, Phys. Rev. D52, 4985(1995).
- [3] G. Barr, T. K. Gaisser and Todor Stanev, Phys. Rev. D39, 3532(1989); V. Agrawal, T. K. Gaisser, Paolo Lipari and Todor Stanev, *ibid.*, D53, 1314(1996)
- [4] T. K. Gaisser, M. Honda, K. Kasahara, H. Lee, S. Midorikawa, V. Naumov, and Todor Stanev, Phys. Rev. D54, 5578(1996).
- [5] Naoya Hata and Paul Langcker, Phys. Rev. D56, 6107 (1997);
- [6] J. N. Bachall, P. I. Krastev and A. Yu. Smirnov, hep-ph/9807216, to be published.
- [7] G. L. Fogli, E. Lisi, A. Marrone, and G. Scioscia, hep-ph/9808205, E. Kh. Akhmedov, A. Dighe, P. Lipari, A. Yu. Smirnov, hep-ph/9808270.
- [8] CHOOZ Collaboration, M. Apollonio *et al.*, Phys. Lett. B420, 397 (1998).
- [9] V. Barger, S. Pakvasa, T. J. Weiler, and K. Whisnant, hep-ph/9806387; Osamu Yasuda, hep-ph/9809205; R. N. Mohapatra and S. Nussinov, hep-ph/9809415, H. Fritzsch and Z. Z. Xing, Phys. Lett. B372 (1996) 265; *ibid* hep-ph/9808272
- [10] Howard Georgi and S. L. Glashow, hep-ph/9808293.